Web of Scie	nce InCites	Journal Citation Reports	Essential Science Indicators	EndNote	Publons	Kopernio	Master Journal Li	st Sign In 🔻	Help 🖣	► English ▼	
Web	o of S	cience							D.	<b>Clarivate</b> Analytics	
Search	Search Re	sults			Тос	ols 👻 Sea	arches and alert	s 👻 Search H	listory	Marked List	
Check full-text	HKBU Library	🕺 Look Up Full Text	Full Text Options 🔻 🕞	Export	Add to	o Marked Lis	it				
										4 2 of 3 ▶	
Opti rang	mally est e, and/o	timating the samp r mid-quartile ran	le mean from the s ge	ample s	ize, me	dian, mi	d-	Citation Ne	twork	<	
By: Luc	By the DH (the Debui) <sup>[1]</sup> : Way V (Way Viage <sup>[2]</sup> : the UN (the timing <sup>[2]</sup> : Tang Tt (Tang Tight) <sup>[1]</sup>							In Web of Science Core Collection			
View Web of Science ResearcherID and ORCID							132	Hig	thly Cited Paper		
STATISTICAL METHODS IN MEDICAL RESEARCH						Times Cited		\$			
Volume	e: 27 Issue: 0 1177/09622	6 Pages: 1785-1805 80216669183								Hot Paper	
Published: JUN 2018 Document Type: Article							Create Citation Alert				
View Journal Impact								All Times Cited Counts			
Abstract							132 in All Databases				
The era of big data is coming, and evidence-based medicine is attracting increasing attention to improve decision making in medical practice via integrating evidence from well designed and conducted clinical research. Meta-analysis						See more counts					
is a statistical technique widely used in evidence-based medicine for analytically combining the findings from											
independent clinical trials to provide an overall estimation of a treatment effectiveness. The sample mean and							21				
standard deviation are two commonly used statistics in meta-analysis but some trials use the median, the minimum						num					
and maximum values, or sometimes the first and third quartiles to report the results. Thus, to pool results in a consistent format, researchers need to transform those information back to the sample mean and standard deviation. In this article, we investigate the optimal estimation of the sample mean for meta-analysis from both theoretical and							Cited References				
						and	View Related Records				

## Most recently cited by:

Jiang, Shi-Qin; Huang, Qiu-Fen; Xie, Wei-Ming; et al.

The association between severe COVID-19 and low platelet count: evidence from 31 observational studies involving 7613 participants. BRITISH JOURNAL OF HAEMATOLOGY

(2020) Shan, Fei; Gao, Chao; Li, Xiao-Long; et al. Short- and Long-Term Outcomes after Laparoscopic Versus Open Gastrectomy for

Elderly Gastric Cancer Patients: A Systematic Review and Meta-analysis. JOURNAL OF LAPAROENDOSCOPIC & ADVANCED SURGICAL TECHNIQUES (2020)

> 26 Since 2013

View All

## **Use in Web of Science**

Web of Science Usage Count

9

Last 180 Days

Learn more

This record is from: Web of Science Core Collection - Science Citation Index Expanded

#### Suggest a correction

If you would like to improve the quality of the data in this record, please suggest a correction.

Journal Information

Performance Trends: Essential Science Indicators Impact Factor: Journal Citation Reports

Reprint Address: Tong, TJ (corresponding author)

🛨 Hong Kong Baptist Univ, Dept Comp Sci, Hong Kong, Hong Kong, Peoples R China.

SAGE PUBLICATIONS LTD, 1 OLIVERS YARD, 55 CITY ROAD, LONDON EC1Y 1SP, ENGLAND

Funding Agency	Show details	Grant Number
НКВИ		FRG2/14-15/077 FRG1/14-15/084 FRG2/15-16/019
Hong Kong Research Gra	HKBU12202114	
HKBU Century Club Spo	nsorship Scheme	

# optimal estimation. Our proposed estimators not only improve the existing ones significantly but also share the same virtue of the simplicity. The real data application indicates that our proposed estimators are capable to serve as rules of thumb and will be widely applied in evidence-based medicine.

### Keywords

Author Keywords: Median; meta-analysis; mid-range; mid-quartile range; optimal weight; sample mean; sample size KeyWords Plus: VITAMIN-D DEFICIENCY; PULMONARY TUBERCULOSIS; METAANALYSIS; CALCIUM

empirical perspectives. A major drawback in the literature is that the sample size, needless to say its importance, is either ignored or used in a stepwise but somewhat arbitrary manner, e.g. the famous method proposed by Hozo etal.

We solve this issue by incorporating the sample size in a smoothly changing weight in the estimators to reach the

#### Author Information

- + Hong Kong Baptist Univ, Dept Math, Hong Kong, Hong Kong, Peoples R China.
- Reprint Address: Wan, X (corresponding author)

# Addresses:

- 🛨 [1] Hong Kong Baptist Univ, Dept Math, Hong Kong, Hong Kong, Peoples R China
- + [2] Hong Kong Baptist Univ, Dept Comp Sci, Hong Kong, Hong Kong, Peoples R China

E-mail Addresses: xwan@comp.hkbu.edu.hk; tongt@hkbu.edu.hk

Funding							
Funding Agency	Show details	Grant Number					
		FRG2/14-15/07					
нкви		FRG1/14-15/08					
		ERG2/15-16/01					

View funding text



Publisher