

A MULTILEVEL SUCCESSIVE ITERATION METHOD FOR NONLINEAR ELLIPTIC PROBLEMS

YUNQING HUANG, ZHONGCI SHI, TAO TANG, AND WEIMIN XUE

ABSTRACT. In this paper, a multilevel successive iteration method for solving nonlinear elliptic problems is proposed by combining a multilevel linearization technique and the cascadic multigrid approach. The error analysis and the complexity analysis for the proposed method are carried out based on the two-grid theory and its multilevel extension. A superconvergence result for the multilevel linearization algorithm is established, which, besides being interesting for its own sake, enables us to obtain the error estimates for the multilevel successive iteration method. The optimal complexity is established for nonlinear elliptic problems in 2-D provided that the number of grid levels is fixed.

1. INTRODUCTION

The multigrid method (MG) has been shown to be one of the most efficient techniques for solving partial differential equations and has been studied by many researchers, see, e.g., Brandt [6] and Hackbusch [12] and the references cited therein. The nested version of the multigrid method or the so-called full multigrid method (FMG) can yield the optimal order of operations $\mathcal{O}(N)$ in obtaining the approximate solution with the accuracy of discretization. In contrast with FMG, Deuffhard [9] proposed a cascadic iteration algorithm which employs nested iterations using the conjugate gradient (CG) method or the preconditioned conjugate gradient (PCG) method instead of using MG at each level. Some adaptive strategies are also proposed, see, e.g., [9, 10]. The efficiency of the cascadic algorithm has been demonstrated numerically in [9, 10], and the comparison is made with the numerical results obtained by using the multilevel preconditioner of Bramble, Pasciak and Xu [5] and the hierarchical preconditioner of Yserentant [27]. The main feature of the cascadic iteration is coarse-grid-correction free, and as a result it can be viewed as a one-way multigrid method. Since the cascadic iteration never goes back to the coarse grids, the error associated with the coarse grids is of relatively low frequencies when the iteration reaches the fine grid, which is very hard to reduce by using a conventional smoother. Hence, the cascadic algorithm has to solve the underlying problems on each level to the same accuracy required for the final level, while FMG solves the underlying problems on each level to the discretization accuracy required for the current level. This implies that the cascadic algorithm may require a large amount of computational time on coarse grids. From the theoretical point of view,

Received by the editor June 6, 2000 and, in revised form, October 12, 2002.

2000 *Mathematics Subject Classification.* Primary 65F10, 65N30, 65N55.

Key words and phrases. Multigrid method, cascadic algorithm, finite element method, nonlinear elliptic problem, error estimate, complexity.

another disadvantage of the cascadic iteration algorithm is that the optimal error estimate can be obtained only under the energy norm.

On the other hand, the main advantage of the cascadic algorithm is its simplicity. It is also efficient for a large class of problems, as demonstrated in [4, 9, 10]. Let the index of the final level be J and that of the current level be j . The iteration number on each grid level can be determined *a priori* or *a posteriori*, which in general depends only on the difference $J - j$, but not on the spatial dimensions. It is the independence of the spatial dimension that yields better efficiency in higher dimensions. In recent years, there have been several analyses and applications of the cascadic iteration algorithm, e.g., Shi and Xu [19, 20, 21] applied the cascadic multigrid technique to elliptic problems with nonconforming elements, to the plate bending problem, and to parabolic problems; and Braess and Dehmen [4] applied the cascadic algorithm to the Stokes equations. On the theoretical side, Shaidurov [18] obtained the optimal complexity in H^2 for the cascadic algorithm with CG as a smoother. Bornemann and Deuffhard [2] analyzed the cascadic algorithm for some general smoothers such as the damped Jacobi, the Gauss-Seidel, etc., under the weaker $H^{1+\alpha}$ -regularity assumption with $0 < \alpha \leq 1$.

The main objective of this paper is to study a multilevel successive iteration algorithm for solving nonlinear elliptic equations. In obtaining the algorithm, a multilevel linearization approach and a cascadic multigrid iteration technique are employed. The error analysis and complexity analysis will be carried out by using the theory for the two-grid method which was first introduced by Xu [24, 25] in approximating nonsymmetric indefinite nonlinear problems. It is based on the facts that the low frequencies are governed by some nonlinear nonsymmetric indefinite operators on the coarse grid and the related high frequencies are governed by some linear symmetric positive definite (SPD) operators on the fine grid. Therefore, we can solve a rather complicated problem on the coarse grids, and then solve an easier problem (linear, SPD) on the fine grid as a correction. If the solution on a coarse grid is sufficiently accurate, then the correction (i.e., the difference between a finer grid solution and the coarse grid solution) can be easily obtained by using simple smoothers. It will be shown that on coarse grids only a fixed number of smoothing iterations are needed. Moreover, the iteration number depends not on the meshsize of the coarse grid but on the number of the refinements used. By extending the two-grid theory to the multigrid case, a multilevel successive iteration algorithm can be proposed to solve a class of nonlinear finite element equations. We note that the idea of using the successive iterations to provide reasonable initial values for linear problems can be traced back to Huang and Liu [14]. However, there had been no theoretical justification until the cascadic multigrid iteration algorithm appeared.

We now state some notation and conventions for later use. Let $\Omega \subset R^d$ be a bounded convex polygonal domain of dimension d , and let $W^{k,p}(\Omega)$ be Sobolev space equipped with the L^p norm and semi-norm:

$$\|u\|_{k,p,\Omega} = \left(\int_{\Omega} \sum_{|\alpha| \leq k} |D^{\alpha} u|^p \right)^{\frac{1}{p}}, \quad |u|_{k,p,\Omega} = \left(\int_{\Omega} \sum_{|\alpha|=k} |D^{\alpha} u|^p \right)^{\frac{1}{p}},$$

where $\alpha = (\alpha_1, \dots, \alpha_d)$ is a multi-index. When $p = 2$ we denote $W^{k,2}(\Omega)$ by $H^k(\Omega)$ and omit the index p in the norm notation. $H_0^1(\Omega)$ consists of functions in $H^1(\Omega)$ that vanish on the boundary $\partial\Omega$. $H^{-1}(\Omega)$ is the dual of $H_0^1(\Omega)$. We shall use the

notation $\lesssim$, $\gtrsim$, $\approx$ as in Xu [26]: when we write $x_1 \lesssim y_1$, $x_2 \gtrsim y_2$, $x_3 \approx y_3$, it means that there exist constants c_1, c_2, c_3, C_3 such that

$$x_1 \leq c_1 y_1, \quad x_2 \geq c_2 y_2, \quad c_3 x_3 \leq y_3 \leq C_3 x_3,$$

where the c_i 's and C_3 are constants independent of the mesh level j and mesh size h_j . These constants may be different at different places. Some special constants will be defined later. Throughout this paper, the Einstein summation convention is used: summation is taken over repeated indices. For example, $a_{ij}b_i$ denotes $\sum_{i=1}^n a_{ij}b_i$.

The rest of the paper is organized as follows. In Section 2 some preliminaries relevant to our error and complexity analysis will be provided. In particular, we will briefly review and study the two-grid method and the multilevel linearization technique. With these preparations, a multilevel successive cascadic iteration algorithm will be proposed and analyzed in Section 3.

2. PRELIMINARIES

Let us consider the following second-order nonlinear elliptic problem:

$$(2.1) \quad \begin{cases} L(u) = -\partial_i(a_i(x, \nabla u)) &= f(x) \quad \text{in } \Omega, \\ u &= 0 \quad \text{on } \partial\Omega. \end{cases}$$

We assume that $a_i(x, y) : \bar{\Omega} \times \mathcal{R}^d \rightarrow \mathcal{R}^1$ is smooth and (2.1) has a unique and nonsingular solution $u \in H_0^1(\Omega) \cap W^{2,d+\epsilon}$ for some $\epsilon > 0$. The linearized operator L' of L at w is defined by the Fréchet derivative as

$$(2.2) \quad L'(w)\phi = -\partial_i(a_{ij}(x, \nabla w)\partial_j\phi),$$

where

$$(2.3) \quad a_{ij}(x, \nabla w) = \frac{\partial a_i(x, \nabla w)}{\partial y_j}.$$

Assume that $a_{ij}(x, \nabla w)$ is SPD for w in a neighborhood of the solution u for (2.1), i.e., there exist two constants α_0 and K such that, $\forall \xi \in \mathcal{R}^d$,

$$(2.4) \quad a_{ij}(x, \nabla w)\xi_i\xi_j \geq \alpha_0|\xi|^2 \quad \forall w \in \mathcal{B}_K,$$

where

$$\mathcal{B}_K = \{w \in W^{1,\infty} : \|w - u\|_{1,\infty} \leq K\}.$$

Let

$$(2.5) \quad A(u, v) := (a_i(x, \nabla u), \partial_i v)$$

and

$$(2.6) \quad A'(w; \phi, v) := (L'(w)\phi, v) = (a_{ij}(x, \nabla w)\partial_j\phi, \partial_i v).$$

Assume that A' is bounded in a neighborhood of u in the following sense:

$$(2.7) \quad |A'(w, \phi, v)| \leq M\|\phi\|_1\|v\|_1 \quad \forall \phi, v \in H_0^1(\Omega), \quad w \in \mathcal{B}_K.$$

Then the weak solution $u \in H_0^1(\Omega)$ for (2.1) is defined by the following equation:

$$(2.8) \quad A(u, v) = (f, v) \quad \forall v \in H_0^1(\Omega),$$

where (f, v) is the standard inner product in Ω . Let $V^h \subset H_0^1(\Omega)$ be a finite element space. The finite element approximation $u_h \in V^h$ for the above problem is then defined by

$$(2.9) \quad A(u_h, v) = (f, v) \quad \forall v \in V^h.$$

For any $u, v, w \in H_0^1$, set $\eta(t) = A(w + t(u - w), v)$. Since

$$\eta(1) - \eta(0) = \eta'(0) + \int_0^1 \eta''(t)(1-t)dt,$$

we obtain the equality

$$(2.10) \quad A(u, v) - A(w, v) = A'(w; u - w, v) + R(w, u, v),$$

where the last term satisfies

$$(2.11) \quad |R(w, u, v)| \lesssim \|u - w\|_{1,2p}^2 \|v\|_{1,p'}, \quad \frac{1}{p} + \frac{1}{p'} = 1.$$

The last estimate is obtained by calculating $\eta''(t)$ directly and by using a Hölder-type inequality, see also [25]. Replacing (w, u) in (2.10) by the solution u of (2.8) and u_h of (2.9) gives

$$(2.12) \quad A'(u; u - u_h, v) = R(u, u_h, v) \quad \forall v \in V^h$$

or

$$(2.13) \quad A'(u_h; u_h - u, v) = R(u_h, u, v) \quad \forall v \in V^h.$$

The existence and uniqueness for the finite element approximation (2.9) and its error estimates can be found in Frehse and Rannacher [11], Rannacher [15], Xu [24], and Chen and Huang [8].

Lemma 2.1 (Xu [24]). *If $u \in W^{2,d+\epsilon}(\Omega)$ and $u_h \in V^h$ are the solution of (2.8) and (2.9) respectively, then the following estimates hold:*

$$(2.14) \quad \begin{aligned} \|u - u_h\|_{1,p} &\lesssim h && \text{if } u \in W^{2,p}, \quad 2 \leq p \leq \infty, \\ \|u - u_h\|_{0,p} &\lesssim h^2 && \text{if } u \in W^{2,p}, \quad 2 \leq p < \infty, \\ \|u - u_h\|_{0,\infty} &\lesssim h^2 |\ln h| && \text{if } u \in W^{2,\infty}. \end{aligned}$$

Suppose that $T^{h_j}, 1 \leq j \leq J$, is a nested quasi-uniform triangulation of Ω and the corresponding linear conforming finite element space is defined by

$$(2.15) \quad V_j = \{v \in C(\overline{\Omega}), v|_e \in P_1, \forall e \in T^{h_j}, v = 0 \text{ on } \partial\Omega\}.$$

Since T^{h_j} is nested, we have

$$V_0 \subset V_1 \subset \cdots \subset V_J \subset H_0^1(\Omega).$$

For simplicity we assume that

$$(2.16) \quad h_j \approx 2^{-j} h_0.$$

The main objective of this work is to propose a multilevel successive iteration method for solving *nonlinear* elliptic equations. Although the problem under investigation is nonlinear, at each fixed level it is linear. Having this in mind, we will carry out some analysis for the two-grid iterative algorithm and the multilevel linearization methods in this section.

2.1. A two-grid analysis. Let us consider the linear model problem to get some intuition:

$$(2.17) \quad (\nabla u, \nabla v) = (f, v) \quad \forall v \in V = H_0^1(\Omega),$$

and the corresponding finite element approximation

$$(2.18) \quad (\nabla u_j, \nabla v) = (f, v) \quad \forall v \in V_j.$$

It is easy to see that

$$(2.19) \quad (\nabla(u_j - u_i), \nabla v) = 0, \quad j \geq i, \quad \forall v \in V_i.$$

This implies that the finite element solution u_i approximating u can also be regarded as a finite element approximation to u_j , $j \geq i$. The increment $u_j - u_i$ is orthogonal to the subspace V_i , indicating that it is of high frequency on the coarser grid T^{h_i} . One then expects to obtain the increment by some simple smoothing iteration.

Now we turn to the nonlinear equation (2.8). Suppose that we have the exact finite element solution u_{h_i} on the level i and want to find an approximation u_j of the finite element solution u_{h_j} on the level j , $j > i$. It follows from (2.10) that if u is the solution of (2.8) and w is an approximation of u , then

$$(2.20) \quad \begin{aligned} A'(w; u, v) &= A'(w; w, v) + A(u, v) - A(w, v) - R(w, u, v) \\ &= A'(w; w, v) + (f, v) - A(w, v) - R(w, u, v). \end{aligned}$$

It is known that $R(w, u, v)$ is a higher order term in $H^1(\Omega)$ and may be neglected. This will lead to the following linearized equation:

$$(2.21) \quad A'(u_{h_i}; u_j, v) = A'(u_{h_i}; u_{h_i}, v) + (f, v) - A(u_{h_i}, v) \quad \forall v \in V_j.$$

The solution u_j of the above problem is an approximation of u_{h_j} . This describes a standard two-grid method of [24]. It is easy to see from (2.21) and (2.10) that

$$(2.22) \quad A'(u_{h_i}; u_j - u_{h_j}, v) = R(u_{h_i}, u_{h_j}, v) \quad \forall v \in V_j,$$

which implies that

$$\|u_j - u_{h_j}\|_{1,2} \lesssim h_i^2.$$

So the discrete accuracy is guaranteed.

Two-grid iteration algorithm.

- (1) Solve the finite element equation (2.9) for u_{h_i} on the level i .
- (2) Set $u_{j,0} = u_{h_i}$, and solve (2.21) by executing m_j smoothing steps on the level j . Let $I_{j,m_j} u_{j,0}$ be the output after m_j steps of iteration: $u_{j,m_j} = I_{j,m_j} u_{j,0}$.

We assume that the error propagation operator $S_{j,m_j} : V_j \rightarrow V_j$ is a linear mapping:

$$(2.23) \quad u_j - I_{j,m_j} u_{j,0} = S_{j,m_j} (u_j - u_{j,0}).$$

We say an iteration is a smoother if it admits the following properties:

$$(2.24) \quad \|S_{j,m_j} v_j\|_a \lesssim \frac{h_j^{-1}}{m_j^\gamma} \|v_j\|_0 \quad \forall v_j \in V_j,$$

$$(2.25) \quad \|S_{j,m_j} v_j\|_a \leq \|v_j\|_a \quad \forall v_j \in V_j,$$

for some constant $0 < \gamma \leq 1$, where $\|\cdot\|_a$ is the a -norm (energy norm) corresponding to the linear system we wish to solve. It is shown that the Gauss-Seidel iteration, SOR, the Richardson iteration, and the damped Jacobi iteration are all smoothers

with the constant $\gamma = \frac{1}{2}$; see, e.g., [12, 26]. Moreover, Shaidurov [18] and Bornemann and Deuffhard [2] proved that the CG method behaves like a linear smoother with $\gamma = 1$.

The following result gives an error bound for the two-grid iteration algorithm.

Assertion 2.1. Let u_j be the exact solution of the linearized equation (2.21) and u_{j,m_j} be an approximation of u_j defined by the above two-grid iteration algorithm; namely, $u_{j,m_j} = I_{j,m_j} u_{j,0}$. If the iteration used is a smoother, then

$$\|u_j - u_{j,m_j}\|_1 \lesssim \frac{2^{2(j-i)}}{m_j^\gamma} h_j,$$

where $0 < \gamma \leq 1$ is defined by (2.24).

Proof. It follows from (2.23)-(2.25) and (2.16) that

$$\begin{aligned} \|u_j - u_{j,m_j}\|_1 &\lesssim \frac{h_j^{-1}}{m_j^\gamma} \|u_j - u_i\|_0 \\ &= \frac{h_j^{-1}}{m_j^\gamma} \|u_j - u + u - u_i\|_0 \\ &\lesssim \frac{h_j^{-1}}{m_j^\gamma} \|u - u_i\|_0 \\ &\lesssim \frac{2^{2(j-i)}}{m_j^\gamma} h_j. \end{aligned}$$

This establishes the desired estimate. $\square$

The above result shows that to achieve the discretization accuracy on the level j it is sufficient to do m_j iterations, with

$$m_j \approx 2^{\frac{2}{\gamma}(j-i)}.$$

We can see that this m_j depends only on the difference $j - i$, and the result is valid for any spatial dimensions.

If we choose $i = j - 1$, then only a fixed number of iterations are needed, i.e., $m \approx 2^{\frac{2}{\gamma}}$, which is independent of the level and the mesh size. However, if we want to iterate the solution on the level i to achieve the discretization accuracy on the level J , then the number of iterations required is $m_i \approx 2^{\frac{2}{\gamma}(J-i)}$.

The generalization of the two-grid algorithm to the multilevel gives the cascadic multigrid iteration scheme. The key idea in the generalization is to use the two-grid technique recursively.

2.2. Multilevel linearization method. Similarly to the projective Newton method [23, 16, 22], we can derive the following multilevel linearization algorithm for solving the nonlinear equation (2.8).

Multilevel linearization algorithm. [1, 13, 24]

- (1) Find $u_0 \in V_0$ such that $A(u_0, v) = (f, v) \quad \forall v \in V_0$.
- (2) For $j = 1, 2, \dots, J$, find $u_j \in V_j$ such that

$$(2.26) \quad A'(u_{j-1}; u_j, v) = A'(u_{j-1}; u_{j-1}, v) + (f, v) - A(u_{j-1}, v) \quad \forall v \in V_j.$$

Let u_{h_j} be the solution of (2.9) in V_j . From (2.26) and (2.10) we have

$$\begin{aligned} A'(u_{j-1}; u_j - u_{j-1}, v) &= (f, v) - A(u_{j-1}, v) \\ &= A(u_{h_j}, v) - A(u_{j-1}, v) \\ &= A'(u_{j-1}; u_{h_j} - u_{j-1}, v) + R(u_{j-1}, u_{h_j}, v), \end{aligned}$$

which gives

$$(2.27) \quad A'(u_{j-1}; u_j - u_{h_j}, v) = R(u_{j-1}, u_{h_j}, v) \quad \forall v \in V_j.$$

We now prove some superconvergence properties for the error between u_{h_j} , the exact solution of (2.9), and u_j , the approximation of u_{h_j} given by (2.26).

Assertion 2.2. Assume $h_0 \ll 1$, and let u be the solution of (2.8). The following results hold:

- If $u \in W^{2,\infty}(\Omega)$, then

$$(2.28) \quad \|u_j - u_{h_j}\|_{1,\infty} \lesssim h_j^2 |\ln h_j|.$$

- If $u \in W^{2,4}(\Omega)$, then

$$(2.29) \quad \|u_j - u_{h_j}\|_{1,2} \lesssim h_j^2.$$

Proof. By Lemma 2.1, we may assume that

$$(2.30) \quad \|u_{h_j} - u_{h_{j-1}}\|_{1,\infty} \leq C_2 h_j.$$

Moreover, we may also choose $h_0 \ll 1$ such that $u_{h_j} \in \mathcal{B}_{K/2}$; namely,

$$(2.31) \quad \|u - u_{h_j}\|_{1,\infty} \leq \frac{K}{2},$$

where the constant K is the same as in (2.4). It is known that the finite element approximation G_h of the regularized Green function of derivative type related to the bilinear form $A'(w; \phi, v)$ satisfies the following inequality [7, 17]:

$$(2.32) \quad \|G_h\|_{1,1} \leq C_0 |\ln h|.$$

Let $C_1 = MC_0/\alpha_0$ and $C = 4C_1C_2^2$. Choose $h_0 \ll 1$ such that

$$2Ch_0 |\ln h_0| \leq C_2.$$

Now we prove (2.28) by induction. For $j = 0$, (2.28) holds since $u_0 = u_{h_0}$. Under the assumptions

$$(2.33) \quad u_{j-1} \in \mathcal{B}_K \quad \text{and} \quad \|u_{j-1} - u_{h_{j-1}}\|_{1,\infty} \leq Ch_{j-1}^2 |\ln h_{j-1}|,$$

we will show that (2.33) is also true when the index $j - 1$ is replaced by j . Taking $v = G_h$ in (2.27), the ellipticity (2.4) and the boundedness (2.7) yield

$$\begin{aligned} \|u_j - u_{h_j}\|_{1,\infty} &\leq C_1 \|u_{h_j} - u_{j-1}\|_{1,\infty}^2 |\ln h_j| \\ &\leq 2C_1 (\|u_{h_j} - u_{h_{j-1}}\|_{1,\infty}^2 + \|u_{h_{j-1}} - u_{j-1}\|_{1,\infty}^2) |\ln h_j| \\ &\leq 2C_1 (C_2^2 h_j^2 + C^2 h_{j-1}^4 |\ln h_{j-1}|^2) |\ln h_j| \\ &\leq 2C_1 (C_2^2 + 4C^2 h_{j-1}^2 |\ln h_{j-1}|^2) h_j^2 |\ln h_j| \\ &\leq 4C_1 C_2^2 h_j^2 |\ln h_j| \\ (2.34) \quad &\leq Ch_j^2 |\ln h_j| \end{aligned}$$

and

$$\begin{aligned}\|u_j - u\|_{1,\infty} &\leq \|u_j - u_{h_j}\|_{1,\infty} + \|u_{h_j} - u\| \\ &\leq Ch_j^2 |\ln h_j| + \frac{K}{2} \leq K.\end{aligned}$$

This completes the induction proof for (2.28). We will now prove (2.29). Let the constant C_2 in (2.30) also be the bounding constant for the $W^{1,4}$ norm, i.e.,

$$(2.35) \quad \|u_{h_j} - u_{h_{j-1}}\|_{1,4} \leq C_2 h_j.$$

Let C_3 be the bounding constant in the inverse estimate of the finite element spaces, i.e.,

$$(2.36) \quad \|v\|_{1,p} \leq C_3 h_j^{\frac{d}{p} - \frac{d}{q}} \|v\|_{1,q} \quad \forall v \in V_j.$$

Assume $h_0 \ll 1$ such that $2CC_3h_0^{1-\frac{d}{4}} \leq C_2$. Again we use induction to prove (2.29). For $j = 0$, (2.29) holds since $u_0 = u_{h_0}$. Assume

$$(2.37) \quad u_{j-1} \in \mathcal{B}_k \quad \text{and} \quad \|u_{j-1} - u_{h_{j-1}}\|_{1,2} \leq Ch_{j-1}^2;$$

we will show that (2.37) is also true when the index $j-1$ is replaced by j . Similarly to the derivation of (2.34), we have

$$\begin{aligned}\|u_j - u_{h_j}\|_{1,2} &\leq C_1 \|u_{h_j} - u_{j-1}\|_{1,4}^2 \\ &\leq 2C_1 (\|u_{h_j} - u_{h_{j-1}}\|_{1,4}^2 + \|u_{h_{j-1}} - u_{j-1}\|_{1,4}^2) \\ &\leq 2C_1 (C_2^2 h_j^2 + C_3^2 h_{j-1}^{-\frac{d}{2}} \|u_{h_{j-1}} - u_{j-1}\|_{1,2}^2) \\ &\leq 2C_1 \left(C_2^2 + 4C^2 C_3^2 h_{j-1}^{2(1-\frac{d}{4})} \right) h_j^2 \\ (2.38) \quad &\leq 4C_1 C_2^2 h_j^2 \leq Ch_j^2,\end{aligned}$$

and

$$\begin{aligned}\|u - u_j\|_{1,\infty} &\leq \|u - u_{h_j}\|_{1,\infty} + C_3 h_j^{-\frac{d}{2}} \|u_{h_j} - u_j\|_{1,2} \\ (2.39) \quad &\leq \frac{K}{2} + C_3 Ch_j^{2-\frac{d}{2}} \leq K.\end{aligned}$$

This completes the proof. $\square$

The superconvergence estimates (2.28) and (2.29) appear to be new. Similar results for the derivatives were obtained by Xu [24] and Bank [1]. The following corollary is a direct consequence of the above assertion.

Corollary 2.1. *Assume $h_0 \ll 1$ and let u be the solution of (2.8).*

• *If $u \in W^{2,\infty}(\Omega)$, then*

$$(2.40) \quad \|u - u_j\|_{1,\infty} \lesssim h_j,$$

$$(2.41) \quad \|u - u_j\|_{0,\infty} \lesssim h_j^2 |\ln h_j|.$$

• *Furthermore, if $u \in W^{2,4}(\Omega)$, then*

$$(2.42) \quad \|u - u_j\|_{1,4} \lesssim h_j,$$

$$(2.43) \quad \|u - u_j\|_{0,2} \lesssim h_j^2.$$

2.3. Inexact multilevel linearization method. In the multilevel linearization method stated above, one must compute u_j exactly by using (2.26) with the initial value u_{j-1} . However, in practice, u_j is only approximated up to some accuracy by using approximation methods or the so-called inexact solvers. In this subsection we analyze the influence of the inexact solver on each grid level to the final grid solution. The error estimates under both the energy norm and the L_2 norm will be obtained. To prove the L_2 norm estimate, a duality argument for nonlinear problems will be developed. We concentrate only on the 2-D case.

Suppose we use an inexact solver in the multilevel linearization method at the level j and obtain the approximation u_j^* , $j = 1, \dots, J$. Let U_j be the exact solution of the linearized equation (2.26) with the initial value u_{j-1}^* , $j \geq 1$. Denote by e_j the error of this solver on the level j , i.e.,

$$(2.44) \quad u_j^* = U_j + e_j.$$

Then the inexact multilevel linearization algorithm is as follows.

Inexact Multilevel Linearization Algorithm.

- (1) Let $U_0 = u_{h_0} \in V_0$ be the solution of $A(U_0, v) = (f, v) \quad \forall v \in V_0$.
- (2) For $j = 1, 2, \dots, J$, find $U_j \in V_j$ such that

$$(2.45) \quad A'(u_{j-1}^*; U_j, v) = A'(u_{j-1}^*; u_{j-1}^*, v) + (f, v) - A(u_{j-1}^*, v) \quad \forall v \in V_j.$$

Let u_{h_j} be the solution of (2.9) in V_j . Similarly to (2.22), we have

$$\begin{aligned} A'(u_{j-1}^*; U_j - u_{j-1}^*, v) &= (f, v) - A(u_{j-1}^*, v) \\ &= A(u_{h_j}, v) - A(u_{j-1}^*, v) \\ &= A'(u_{j-1}^*; u_{h_j} - u_{j-1}^*, v) + R(u_{j-1}^*, u_{h_j}, v). \end{aligned}$$

Hence

$$(2.46) \quad A'(u_{j-1}^*; U_j - u_{h_j}, v) = R(u_{j-1}^*, u_{h_j}, v) \quad \forall v \in V_j.$$

Now we turn to the error analysis for the solution U_j of (2.45) given by the inexact multilevel linearization algorithm and the exact nonlinear solution u_{h_j} on the level j .

Assertion 2.3. Assume $h_0 \ll 1$, and let u be the solution of (2.8). If $u \in W^{2,4}(\Omega)$ and the error e_j , defined by (2.44), satisfies

$$\|e_j\|_1 \leq \delta h_j, \quad \|e_j\|_1 \leq C_e \|e_{j+1}\|_1, \quad j = 0, 1, \dots, J-1,$$

then

$$(2.47) \quad \|U_j - u_{h_j}\|_{1,2} \lesssim h_j^2 + \|e_j\|_1, \quad j \leq J.$$

Furthermore, if Ω is smooth or convex, then

$$(2.48) \quad \|U_j - u_{h_j}\|_{0,2} \lesssim h_j^2 + \|e_j\|_1^2 |\ln h_j|^{\frac{1}{2}}, \quad j \leq J.$$

Proof. We prove the above results by induction again. If u_{j-1}^* is in the neighborhood of u , using (2.11) and (2.46), we have

$$\begin{aligned} \|U_j - u_{h_j}\|_{1,2} &\leq C_1 \|u_{h_j} - u_{j-1}^*\|_{1,4}^2 \\ &\leq 2C_1 (\|u_{h_j} - u_{h_{j-1}}\|_{1,4}^2 + \|u_{h_{j-1}} - u_{j-1}^*\|_{1,4}^2) \\ (2.49) \quad &\leq 2C_1 \{C_2^2 h_j^2 + 2C_3^2 h_{j-1}^{-1} (\|u_{h_{j-1}} - U_{j-1}\|_{1,2}^2 + \|e_{j-1}\|_{1,2}^2)\}, \end{aligned}$$

where C_2 and C_3 are constants defined in (2.35) and (2.36), respectively. Let $C = 4C_1C_2^2$. Suppose h_0 is small enough so that

$$(2.50) \quad \|u - u_{h_j}\|_{1,\infty} \leq K/2, \quad 64C_1C_3^2h_0 \leq 1, \quad CC_3h_0 \leq K/4.$$

Suppose $\delta \ll 1$ is such that

$$(2.51) \quad C_3C_e\delta \leq K/4, \quad 12C_1C_3^2C_e^2\delta \leq 1.$$

For $j = 0$ we have $U_0 - u_{h_0} = 0$, so the estimate (2.47) is valid. Furthermore, by (2.50) and (2.51) we have

$$(2.52) \quad \begin{aligned} \|u - u_0^*\|_{1,\infty} &\leq \|u - u_{h_0}\|_{1,\infty} + \|e_0\|_{1,\infty} \\ &\leq K/2 + C_3\delta \leq K, \end{aligned}$$

which implies that $u_0^* \in \mathcal{B}_K$. Suppose (2.47) holds for $j \leq i-1$, i.e.,

$$(2.53) \quad \|U_j - u_{h_j}\|_1 \leq Ch_j^2 + \|e_j\|_1^2, \quad j < i.$$

It can be verified that $u_{i-1}^* \in \mathcal{B}_K$, and

$$(2.54) \quad \begin{aligned} \|u - u_{i-1}^*\|_{1,\infty} &\leq \|u - U_{i-1}\|_{1,\infty} + \|e_{i-1}\|_{1,\infty} \\ &\leq K/2 + C_3C_e\delta \leq K. \end{aligned}$$

The assumption (2.53), together with (2.49)-(2.51), yields

$$(2.55) \quad \begin{aligned} \|U_i - u_{h_i}\|_{1,2} &\leq 2C_1(C_2^2 + 16CC_3^2h_i)h_i^2 + 12C_1C_3^2C_e^2\delta\|e_i\|_1 \\ &\leq Ch_i^2 + \|e_i\|_1, \end{aligned}$$

which means that (2.47) is valid for $j = i$. It remains to verify that U_i is still in the neighborhood of u , i.e., $U_i \in \mathcal{B}_K$, in order to complete the proof of (2.47). In fact, using (2.50), (2.51) and the inverse estimates gives

$$(2.56) \quad \begin{aligned} \|u - U_i\|_{1,\infty} &\leq \|u - u_{h_i}\|_{1,\infty} + C_3h_i^{-1}\|u_{h_i} - U_i\|_1 \\ &\leq \frac{K}{2} + C_3(Ch_i + \delta) \leq K. \end{aligned}$$

Finally, we prove the optimal L_2 error estimate for u_j generated by the inexact multilevel linearization algorithm. We use a duality argument. Construct the following auxiliary problem: Find $w \in H_0^1(\Omega)$ such that

$$(2.57) \quad A'(u_{j-1}^*; v, w) = (v, U_j - u_{h_j}) \quad \forall v \in H_0^1(\Omega).$$

Let $w_j \in V_j$ be the finite element approximation of w in V_j , and set $v = U_j - u_{h_j}$ in (2.57). It follows from (2.46), (2.11) and (2.7) that

$$(2.58) \quad \begin{aligned} \|U_j - u_{h_j}\|_{0,2}^2 &= A'(u_{j-1}^*; U_j - u_{h_j}, w) \\ &= A'(u_{j-1}^*; U_j - u_{h_j}, w - w_j) + A'(u_{j-1}^*; U_j - u_{h_j}, w_j) \\ &= A'(u_{j-1}^*; U_j - u_{h_j}, w - w_j) + R(u_{j-1}^*, u_{h_j}, w_j) \\ &\lesssim \|U_j - u_{h_j}\|_1 \|w - w_j\|_1 + \|u_{h_{j-1}} - u_{h_j}\|_{1,4}^2 \|w_j\|_1 \\ &\quad + (\|U_{j-1} - u_{h_{j-1}}\|_1^2 + \|e_{j-1}\|_1^2) \|w_j\|_{1,\infty} \\ &\lesssim (h_j^2 + \|e_j\|_1^2 |\ln h_j|^{\frac{1}{2}}) \|w\|_2. \end{aligned}$$

In the last inequality we have used the fact that $\|w_j\|_{1,\infty} \lesssim |\ln h_j|^{\frac{1}{2}} \|w\|_2$, which can be obtained by using the inverse estimates, the $W^{1,p}$ -stability property and the embedding theorem. Hence, applying standard a priori estimates yields (2.48). $\square$

3. MULTILEVEL SUCCESSIVE ITERATION METHOD

Based on the preparations in the last section, we propose the following multilevel successive iterative algorithm for solving the nonlinear equation (2.9).

Multilevel Successive Iteration Algorithm.

- (1) Solve the nonlinear problem $A(u_{h_0}, v) = (f, v) \quad \forall v \in V_0$ by an appropriate solver to obtain u_0^* with the accuracy δh_0 and h_0^2 in the H^1 -norm and L_2 -norm, respectively.
- (2) For $j = 1, 2, \dots, J$, let I_{j,m_j} denote m_j basic smoothing iterations on the level j with initial data u_{j-1}^* for solving the linearized equation

$$A'(u_{j-1}^*; U_j, v) = A'(u_{j-1}^*; u_{j-1}^*, v) + (f, v) - A(u_{j-1}^*, v) \quad \forall v \in V_j,$$

$$\text{and set } u_j^* = I_{j,m_j} u_{j-1}^*.$$

Now we give the error analysis for the exact solution U_j and its approximation u_j^* in the above algorithm. Let $u_j^* = U_j - e_j$. Then

$$\begin{aligned} e_j &= U_j - u_j^* = S_{j,m_j}(U_j - u_{j-1}^*) \\ (3.1) \quad &= S_{j,m_j}(U_j - U_{j-1}) + S_{j,m_j}e_{j-1}. \end{aligned}$$

Define the level-dependent norm

$$(3.2) \quad \|v\|_{a,j}^2 = A'(u_{j-1}^*; v, v) \quad \forall v \in V_j, \quad j \geq 1.$$

Lemma 3.1. Assume $h_0 \ll 1$, and let u be the solution of (2.8). If $u \in W^{2,4}(\Omega)$ and the error e_j , defined by (2.44), satisfies

$$\|e_j\|_1 \leq \delta h_j, \quad \|e_j\|_1 \leq C_e \|e_{j+1}\|_1, \quad j = 0, 1, \dots, J-1,$$

then for any $v \in V_j$, $j \geq 2$,

$$(3.3) \quad \|v\|_{a,j}^2 \leq \left(1 + Ch_j^{\frac{1}{2}} + C\|e_{j-1}\|_{1,\infty} + C\|e_{j-2}\|_{1,\infty}\right) \|v\|_{a,j-1}^2,$$

where C is a constant independent of the mesh size h and the level j .

Proof. It follows from the definitions (2.6) and (3.2) that

$$\begin{aligned} \|v\|_{a,j}^2 &= A'(u_{j-1}^*; v, v) \\ &= \|v\|_{a,j-1}^2 + A'(u_{j-1}^*; v, v) - A'(u_{j-2}^*; v, v) \\ (3.4) \quad &\leq \|v\|_{a,j-1}^2 + C|u_{j-1}^* - u_{j-2}^*|_{1,\infty} \|v\|_1^2. \end{aligned}$$

Using the inverse estimate (2.36) and Lemma 2.1, we obtain

$$\begin{aligned} |u_{h_{j-1}} - u_{h_{j-2}}|_{1,\infty} &\lesssim h_{j-1}^{-\frac{1}{2}} \|u_{h_{j-1}} - u_{h_{j-2}}\|_{1,4} \\ (3.5) \quad &\lesssim h_j^{\frac{1}{2}}. \end{aligned}$$

If we use estimates (3.4) and (3.5), then Assertion 2.3 together with the identities

$$\begin{aligned} u_{j-1}^* - u_{j-2}^* &= U_{j-1} - U_{j-2} - e_{j-1} - e_{j-2}, \\ U_{j-1} - U_{j-2} &= (U_{j-1} - u_{h_{j-1}}) + (u_{h_{j-1}} - u_{h_{j-2}}) + (u_{h_{j-2}} - U_{j-2}), \end{aligned}$$

yields the desired inequality (3.3). $\square$

Theorem 3.1. Assume that $d = 2, h_0 \ll 1$, $u \in W^{2,4}(\Omega)$ and Ω is smooth or convex. If we choose the number of iterations as $m_j = m\tilde{m}_j$ with

$$(3.6) \quad \tilde{m}_j = \begin{cases} [J^2 2^{2(J-j)}] + 1 & \text{if } \gamma = \frac{1}{2}, \\ [J^{\frac{1}{2}} 2^{\beta(J-j)}] + 1, 1 < \beta < 2, & \text{if } \gamma = 1, \end{cases}$$

where m is a proper constant, then the multilevel successive iteration algorithm admits the error bound

$$(3.7) \quad \|e_j\|_{a,j} \lesssim h_J, \quad j = 1, 2, \dots, J,$$

and the total number of operations is bounded by

$$(3.8) \quad W_J \lesssim \begin{cases} J^3 N_J, & \gamma = \frac{1}{2}, \\ \sqrt{J} N_J, & \gamma = 1. \end{cases}$$

Proof. Without loss of generality, we may choose $h_0 \ll 1$ as in (2.31) such that

$$(3.9) \quad \|u - U_j\|_{1,\infty} \leq K/2.$$

We establish the error bound (3.7) by induction. For $j = 0$, the result is obvious. Suppose that

$$(3.10) \quad \|e_j\|_a < h_J K/2$$

for $j \leq i - 1$. Then

$$(3.11) \quad \|u_j^* - u\|_{1,\infty} \leq K, \quad j = 1, \dots, i - 1,$$

$$(3.12) \quad \|e_j\|_{1,\infty} \lesssim h_j^{-1} \|e_j\|_{1,2} \lesssim \frac{h_J}{h_j} \lesssim 2^{j-J}, \quad j = 1, \dots, i - 1.$$

By using Lemma 3.1, the smoothing property (2.25), and (3.12), we have

$$\begin{aligned} \|e_i\|_{a,i} &\leq \|S_{i,m_i}(U_i - U_{i-1})\|_{a,i} + \|S_{i,m_i}e_{i-1}\|_{a,i} \\ &\leq \|S_{i,m_i}(U_i - U_{i-1})\|_{a,i} + \|e_{i-1}\|_{a,i} \\ &\leq \|S_{i,m_i}(U_i - U_{i-1})\|_{a,i} + \left(1 + C2^{j-1-J} + Ch_{i-1}^{\frac{1}{2}}\right) \|e_{i-1}\|_{a,i-1} \\ &\leq \sum_{j=1}^i \prod_{l=j}^{i-1} \left(1 + C2^{l-J} + Ch_l^{\frac{1}{2}}\right) \|S_{j,m_j}(U_j - U_{j-1})\|_{a,j} \\ &\leq \tilde{C}_1 \sum_{j=1}^i \|S_{j,m_j}(U_j - U_{j-1})\|_{a,j}. \end{aligned}$$

Furthermore, it follows from the smoothing property (2.24) and the superconvergence result (2.41) that

$$(3.13) \quad \begin{aligned} \|e_i\|_{a,i} &\leq \frac{\tilde{C}_1}{m^\gamma} \sum_{j=1}^i \left(\frac{h_j}{\tilde{m}_j^\gamma} + \frac{h_j^{-1} h_J^2 |\ln h_j|^{\frac{1}{2}}}{\tilde{m}_j^\gamma} \right) \\ &\leq \frac{\tilde{C}_1 \tilde{C}_2}{m^\gamma} h_J \end{aligned}$$

with

$$\tilde{C}_2 = \left(\sum_{j=1}^J \frac{2^{J-j}}{\tilde{m}_j^\gamma} + \sum_{j=1}^J \frac{J^{\frac{1}{2}} 2^{j-J}}{\tilde{m}_j^\gamma} \right),$$

where the iteration number $\tilde{m}_j$ is given by (3.6). Hence

$$\|e_i\|_{a,i} \leq \frac{K}{2} h_J$$

provided m is sufficiently large. So (3.7) holds for $j = i$, as long as u_i^* is still in the neighborhood of u . The latter requirement can be verified by the following observations:

$$\begin{aligned} \|u_i^* - u\|_{1,\infty} &\leq \|u_i^* - U_i\|_{1,\infty} + \|U_i - u\|_{1,\infty} \\ (3.14) \quad &\leq \frac{K}{2} + \frac{K}{2} \leq K. \end{aligned}$$

Therefore, (3.7) is proved. Finally, we will estimate the total number of operations (3.8). For $\gamma = 1/2$,

$$\begin{aligned} W_J &= \sum_{j=1}^J m_j N_j \lesssim \sum_{j=1}^J J^2 2^{2(J-j)} 2^{2j} N_0 \\ (3.15) \quad &\lesssim J^3 N_J; \end{aligned}$$

while for $\gamma = 1$,

$$\begin{aligned} W_J &= \sum_{j=1}^J m_j N_j \lesssim \sum_{j=1}^J J^{\frac{1}{2}} 2^{\beta(J-j)} 2^{2j} N_0 \\ &\lesssim J^{\frac{1}{2}} N_J \sum_{j=1}^J 2^{(\beta-2)(J-j)} \\ (3.16) \quad &\lesssim J^{\frac{1}{2}} N_J. \end{aligned}$$

This completes the proof of this theorem. $\square$

Theorem 3.1 shows that for a fixed grid level J the complexity of the proposed multilevel successive iteration algorithm is optimal, and is proportional to the total number of unknowns. For an arbitrary grid level the algorithm is quasi-optimal, since $J \approx \ln N_J$ asymptotically.

We close this section with the following observations:

- First, we only considered the error estimate in the H^1 -norm in this paper. However, it is easy to see that for a fixed level J , we can obtain the optimal convergence rate in any norm, provided that the operation number is proportional to the number of unknowns on the final grid. On the other hand, if the mesh level J is arbitrary as in the case of cascadic iterations, then only the H^1 -norm estimate can be kept optimal.
- Although our analysis for nonlinear equations is given only for the 2-D case, it can be extended to 3-D problems when the mesh level J is fixed. However, if the mesh level J is arbitrary, the situation is more complicated. In this case, the cascadic multigrid method only achieves the optimal accuracy in the energy norm, and as a result the most difficult step is to verify whether the iterative approximation u_j^* remains in the neighborhood of u uniformly in the $W^{1,\infty}$ -norm. Unfortunately, this is not the case for strongly nonlinear equations in 3-D. Therefore, it seems difficult to extend the 2-D analysis presented in this section to deal with 3-D problems when J is not fixed. This remains one of our further research topics.

ACKNOWLEDGMENTS

We would like to thank an anonymous referee for helpful suggestions. The research of Y. Huang and Z. Shi was supported in part by special funds for Major State Basic Research Projects of China, and Y. Huang was also supported in part by the National Education Ministry. The research of T. Tang and W. Xue was supported by the FRG grants of the Hong Kong Baptist University and the RGC grants of the Hong Kong Research Grants Council.

REFERENCES

- [1] R. BANK, *Analysis of a multilevel iterative method for nonlinear finite element equations*. Math. Comp., 39 (1982), 453-465. MR **83j**:65105
- [2] F. BORNEMANN AND P. DEUFLHARD, *The cascadic multigrid method for elliptic problems*, Numer. Math., 75 (1996), 135-152. MR **98a**:65175
- [3] F. BORNEMANN AND P. DEUFLHARD, *The cascadic multigrid method*. Proceedings of the 8th international conference on domain decomposition methods, R.Glowinski, J.Periaux, Z. Shi and O.Widlund eds., Wiley and Sons, 1997, pp. 205-212.
- [4] D. BRAESS AND W. DEHMEN, *A cascade algorithm for the Stokes equations*. Numer. Math., 82 (1999), 179-191. MR **2000c**:65114
- [5] J.H. BRAMBLE, J.E. PASCIAK AND J. XU, *Parallel multilevel preconditioner*. Math. Comp., 55 (1990), 1-22. MR **90k**:65170
- [6] A. BRANDT, *Multilevel adaptive solutions to boundary-value problems*. Math. Comp., 31 (1977), 333-390. MR **55**:4714
- [7] C. CHEN AND Y. HUANG, *$W^{1,p}$ -Stability of finite element method for elliptic problems*. Hunan Annals of Mathematics, 6 (1986), 81-89. (Chinese) MR **92e**:65144
- [8] C. CHEN AND Y. HUANG, *High accuracy theory of finite element methods*, Hunan Science and Technology Press, 1995. (Chinese)
- [9] P. DEUFLHARD, *Cascadic conjugate gradient methods for elliptic partial differential equations*. Proceedings of the 7th international conference on domain decomposition methods, eds D. Keyes and J. Xu, AMS, Providence, 1994, pp. 29-42. MR **95i**:65006
- [10] P. DEUFLHARD, P. LEINEN AND H. YSERENTANT, *Concepts of an adaptive hierarchical finite element code*. IMPACT 1 (1989), 3-35.
- [11] J. FREHSE AND R. RANNACHER, *Asymptotic L_∞ -error estimate for linear finite element approximation of quasilinear boundary value problems*. SIAM J. Numer. Anal., 15 (1978), 418-431. MR **58**:19224
- [12] W. HACKBUSCH, *Multi-grid methods and applications*. Springer Verlag, Berlin-Heidelberg-New York, 1992. MR **87e**:65082
- [13] Y. HUANG AND Y. CHEN, *A multilevel iterate correction method for solving nonlinear singular problems*, Natural Science J of Xiangtan Univ. 16 (1994), 23-26. (Chinese)
- [14] H. HUANG AND G. LIU, *Successive refinement method for solving elliptic boundary value problems*. Mathematica Numerica Sinica, 2 (1978), 41-52, and 3 (1978), 28-35. (Chinese)
- [15] R. RANNACHER, *On the finite element approximation of general boundary value problems in nonlinear elasticity*. Calcolo, 17 (1980), 175-193. MR **82f**:73019
- [16] R. RANNACHER, *On the convergence of the Newton-Raphson method for strongly nonlinear finite element equations*. In Nonlinear Computational Mechanics, eds P. Wriggers and W. Wagner, Springer Verlag, 1991.
- [17] R. RANNACHER AND R. SCOT, *Some optimal error estimates for piecewise linear finite element approximations*. Math. Comp., 38 (1982), 437-445. MR **83e**:65180
- [18] V. SHADUROV, *Some estimates of the rate of convergence for the cascadic conjugate gradient method*. Comp. Math. Appl., 31 (1996), 161-171.
- [19] Z. SHI AND X. XU, *Cascadic multigrid method for elliptic problems*, East-West J. Numer. Math., 7 (1999), 199-222. MR **2000h**:66185
- [20] Z. SHI AND X. XU, *Cascadic multigrid method for a plate bending problem*. East-West J. Numer. Math., 6 (1998), 137-153. MR **99g**:73109
- [21] Z. SHI AND X. XU, *Cascadic multigrid method for parabolic problem*. J. Comput. Math., 18 (2000), 551-560. MR **2001e**:65151

- [22] G. VAINIKKO, *Galerkin's perturbation method and the general theory of approximation for nonlinear equations*. USSR Comput. Math. and Math. Phys., 7 (1967), 1-41. MR **36**:1095
- [23] K. WITSCH, *Projective Newton-Verfahren und Adwendungen auf nicht linear Randwertaufgaben*. Numer. Math., 31 (1978/1979), 209-230. MR **80a**:65160
- [24] J. XU, *Two-grid discretization techniques for linear and nonlinear PDEs*. SIAM J. Numer. Anal., 33, (1996), 1759-1777. MR **97i**:65169
- [25] J. XU, *A novel two-grid method for semilinear equations*. SIAM J. Sci. Comput., 15 (1994), 231-237. MR **94m**:65178
- [26] J. XU, *Iterative methods by space decomposition and subspace correction*, SIAM Review, 34 (1992), 581-613. MR **93k**:65029
- [27] H. YSERENTANT, *On the multilevel splitting of finite element spaces*. Numer. Math., 49 (1986), 379-412. MR **88d**:65068a; MR **88d**:65068b

DEPARTMENT OF MATHEMATICS AND INSTITUTE FOR COMPUTATIONAL AND APPLIED MATHEMATICS, XIANGTAN UNIVERSITY, XIANGTAN, HUNAN 411105, PEOPLES REPUBLIC OF CHINA
E-mail address: `huangyq@xtu.edu.cn`

INSTITUTE OF COMPUTATIONAL MATHEMATICS, CHINESE ACADEMY OF SCIENCES, BEIJING, 100080, PEOPLES REPUBLIC OF CHINA
E-mail address: `shi@lsec.cc.ac.cn`

DEPARTMENT OF MATHEMATICS, HONG KONG BAPTIST UNIVERSITY, KOWLOON TONG, HONG KONG
E-mail address: `ttang@math.hkbu.edu.hk`

DEPARTMENT OF MATHEMATICS, HONG KONG BAPTIST UNIVERSITY, KOWLOON TONG, HONG KONG
E-mail address: `wmxue@math.hkbu.edu.hk`