The Sharpness of Kuznetsov's $O(\sqrt{\Delta x})L^1$-Error Estimate for Monotone Difference Schemes

Tao Tang, Zhen-Huan Teng

Mathematics of Computation, Volume 64, Issue 210 (Apr., 1995), 581-589.

Your use of the JSTOR database indicates your acceptance of JSTOR's Terms and Conditions of Use. A copy of JSTOR's Terms and Conditions of Use is available at http://www.jstor.org/about/terms.html, by contacting JSTOR at jstor-info@umich.edu, or by calling JSTOR at (888)388-3574, (734)998-9101 or (FAX) (734)998-9113. No part of a JSTOR transmission may be copied, downloaded, stored, further transmitted, transferred, distributed, altered, or otherwise used, in any form or by any means, except: (1) one stored electronic and one paper copy of any article solely for your personal, non-commercial use, or (2) with prior written permission of JSTOR and the publisher of the article or other text.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

Mathematics of Computation is published by American Mathematical Society. Please contact the publisher for further permissions regarding the use of this work. Publisher contact information may be obtained at http://www.jstor.org/journals/ams.html.

Mathematics of Computation
©1995 American Mathematical Society

JSTOR and the JSTOR logo are trademarks of JSTOR, and are Registered in the U.S. Patent and Trademark Office. For more information on JSTOR contact jstor-info@umich.edu.

©2001 JSTOR

http://www.jstor.org/
Tue Jan 2 02:32:58 2001

MATHEMATICS OF COMPUTATION
VOLUME 64, NUMBER 210
APRIL 1995, PAGES 581–589

THE SHARPNESS OF KUZNETSOV'S $O(\sqrt{\Delta x})$ L^1-ERROR ESTIMATE FOR MONOTONE DIFFERENCE SCHEMES

TAO TANG AND ZHEN-HUAN TENG

ABSTRACT. We derive a lower error bound for monotone difference schemes to the solution of the linear advection equation with BV initial data. A rigorous analysis shows that for any monotone difference scheme the lower L^1-error bound is $O(\sqrt{\Delta x})$, where Δx is the spatial stepsize.

1. INTRODUCTION

Conservative monotone difference schemes, which include the Lax-Friedrichs scheme, Godunov's scheme, and the Engquist-Osher scheme [3], play an important role in both theoretical analysis and practical computation for hyperbolic conservation laws. From the viewpoint of numerical computation, accuracy and error bounds are of particular interest. Harten, Hyman, and Lax [4] pointed out that the monotone difference schemes are of at most first-order accuracy and Kuznetsov [6] showed that their (upper) L^1-error bound is $O(\sqrt{\Delta x})$ as Δx goes to zero, where Δx is the spatial stepsize.

In this paper we demonstrate that *all* monotone schemes applied to linear first-order conservation laws in one dimension have a best possible $\sqrt{\Delta x}$ rate of convergence when applied to discontinuous data.

A $(p+q+1)$-point conservative finite difference scheme

$$\begin{aligned} v_j^{n+1} &= H(v_{j-p}^n, v_{j-p+1}^n, \dots, v_{j+q}^n) \\ &= v_j^n - \lambda[\bar{f}(v_{j-p+1}^n, \dots, v_{j+q}^n) - \bar{f}(v_{j-p}^n, \dots, v_{j+q-1}^n)] \end{aligned} \tag{1.1}$$

is said to be monotone if H is a monotone nondecreasing function of each of its arguments, and is said to be consistent with a scalar conservation law

$$\frac{\partial u}{\partial t} + \frac{\partial f(u)}{\partial x} = 0, \qquad x \in \mathbb{R}, \quad t > 0, \tag{1.2a}$$

$$u|_{t=0} = u_0(x), \tag{1.2b}$$

if the numerical flux $\bar{f}$ satisfies

$$\bar{f}(w, \dots, w) = f(w), \tag{1.3}$$

Received by the editor August 23, 1993 and, in revised form, March 7, 1994.
1991 *Mathematics Subject Classification.* Primary 65M25; Secondary 76D05.
Key words and phrases. Monotone difference scheme, error estimate, lower error bound.
The first author was supported by NSERC Canada (Grants OGP0105545 and BEF0141031) and also partially supported by the National Natural Science Foundation of China.
The second author was supported by the National Natural Science Foundation of China and the Science Fund of the Education Commission of China.

©1995 American Mathematical Society
0025-5718/95 $1.00 + $.25 per page

where $\lambda = \Delta t/\Delta x = \text{const}$, p and q are given nonnegative integers, and

$$v_j^0 = T_{\Delta x}(u_0)(x_j) = \frac{1}{\Delta x}\int_{x_j-\Delta x/2}^{x_j+\Delta x/2} u_0(x)\,dx, \quad x_j = j\Delta x. \tag{1.4}$$

Stability, convergence, and error estimates for monotone difference schemes can be found in [2], [6], and [8].

It is easy to see that if (1.2) is the linear advection equation

$$\frac{\partial u}{\partial t} + a\frac{\partial u}{\partial x} = 0 \qquad (a = \text{const}), \tag{1.5a}$$

$$u|_{t=0} = u_0(x), \tag{1.5b}$$

then a linear $(p+q+1)$-point monotone difference scheme is of the form

$$v_j^{n+1} = \sum_{s=-p}^{q} a_s v_{j+s}^n, \tag{1.6}$$

where

$$a_s \geq 0 \quad \text{for} \quad s = -p, \ldots, q. \tag{1.7}$$

The consistency condition (1.3) implies that

$$\sum_{s=-p}^{q} a_s = 1 \tag{1.8}$$

and

$$\sum_{s=-p}^{q} s a_s = -\lambda a. \tag{1.9}$$

Denote

$$\mathscr{S}_1 = \{s \mid a_s > 0\} \quad \text{and} \quad \mathscr{S}_1 \backslash s_0 = \{s \mid s \in \mathscr{S}_1 \text{ and } s \neq s_0\},$$

where s_0 is an index which satisfies $a_{s_0} = \max_{s\in\mathscr{S}_1} a_s$. For the analysis of (1.6), we introduce

$$v_{\Delta x}(x, t) = v_j^n \quad \text{for} \quad (x, t) \in [x_{j-1/2}, x_{j+1/2}) \times [t_n, t_{n+1}), \tag{1.10}$$

where $x_{j+1/2} = (j+1/2)\Delta x$, $t_n = n\Delta t$, $n \in \mathbb{N}$ and $j \in \mathbb{Z}$.

In this paper we will prove the following theorem.

Theorem. *Any monotone difference scheme* (1.6), *which is consistent with* (1.5), *has the following* L^1*-error bounds: for any* $M > 0$ *and* $t > 0$

$$\begin{aligned} c(p, q)\, M \sum_{s\neq s_0} \sqrt{a_s}\sqrt{\frac{t}{\lambda}}\sqrt{\Delta x} &\leq \sup_{|u_0|_{BV}\leq M} \|v_{\Delta x}(\cdot, t) - u(\cdot, t)\|_{L^1(\mathbb{R})} \\ &\leq M\left[2\sqrt{\sum_s s^2 a_s - \lambda^2 a^2}\sqrt{\frac{t}{\lambda}}\sqrt{\Delta x} + \Delta x\right], \end{aligned} \tag{1.11}$$

provided that Δx *is small enough. Here,* $c(p, q) > 0$ *is a constant depending only on* p *and* q, $u(x, t)$ *is the solution of* (1.5) *and*

$$|u_0|_{BV} = \sup_{|h|\neq 0} \frac{1}{|h|}\|u_0(\cdot + h) - u_0(\cdot)\|_{L^1(\mathbb{R})}. \tag{1.12}$$

Remark 1. The lower bound of (1.11) indicates that, except in a trivial case ($a_s = 0$ for $s \neq s_0$, a pure translation), any monotone difference scheme applied to a linear advection equation has an L^1 convergence order of at most one-half in the class BV of solutions.

Remark 2. Several authors have studied error estimates for difference schemes to first-order hyperbolic equations by using Fourier methods (see [1,5] and references therein). However, to the best of our knowledge, none of these includes a lower error bound for monotone difference schemes in the presence of discontinuous initial data.

2. Some lemmas

A key step in proving the lower error bound of (1.11) is to get a precise lower bound of a sum of terms with multi-indices running over a set J_n (see the right-hand side of (3.7)). Lemma 2 and Lemma 4 below provide a precise lower cardinality of J_n and precise lower bounds of the summand terms, respectively. Consequently, they yield the desired lower bound. Lemma 1 gives a multinomial equality, while Lemma 3 is a generalized de Moivre theorem [7], which gives an asymptotic formula for the multinomial probabilities.

Lemma 1. *If* $\mathbf{a} = (a_{-p}, \dots, a_q) \in \mathbb{R}^{p+q+1}$ *satisfies* (1.7) *and* (1.8), *then*

$$\sum_{|\alpha|=n} C_n(\alpha)\, \mathbf{a}^\alpha = 1, \tag{2.1}$$

where $\alpha = (\alpha_{-p}, \dots, \alpha_q) \in \mathbb{N}^{p+q+1}$, $|\alpha| = \alpha_{-p} + \cdots + \alpha_q$, $\mathbf{a}^\alpha = a_{-p}^{\alpha_{-p}} \cdots a_q^{\alpha_q}$ *and* $C_n(\alpha)$ *is the multinomial coefficient defined by*

$$C_n(\alpha) = \frac{n!}{\alpha_{-p}! \cdots \alpha_q!}. \tag{2.2}$$

Proof. The above equality can be easily derived, and the proof is omitted. □

Denote

$$I_n = \{\alpha \,|\, |\alpha| = n, \quad \alpha \in \mathbb{N}^{p+q+1}\}, \tag{2.3}$$

(2.4a)

$$\mathbb{J}_n = \Big\{\xi \;\Big|\; \xi \in \mathbb{R}^{p+q+1},\ \xi_s = a_s n + y_s\sqrt{n}\ (s \in \mathscr{S}_1) \text{ and } \xi_s = 0\ (s \notin \mathscr{S}_1);\ y_s \in [\bar{y}_s,\ \bar{y}_s + \Delta y_s]\ (s \in \mathscr{S}_1 \backslash s_0) \text{ and } \sum_{s\in\mathscr{S}_1} y_s = 0\Big\}$$

and

$$J_n = \{\alpha \,|\, \alpha \in \mathbb{J}_n \cap \mathbb{N}^{p+q+1}\}, \tag{2.4b}$$

where $\mathbf{a} = (a_{-p}, \dots, a_q)$ satisfies (1.7) and (1.8).

Lemma 2. *For sufficiently large n,*

$$|J_n| \geq |\mathscr{S}_1|/2 \Big(\prod_{s\in\mathscr{S}_1\backslash s_0} |\Delta y_s|\Big) n^{(|\mathscr{S}_1|-1)/2}, \tag{2.5}$$

where $|J_n|$ *and* $|\mathscr{S}_1|$ *are the cardinalities of* J_n *and* $\mathscr{S}_1$, *respectively.*

Proof. Since the set J_n consists of all lattice points of $\mathbb{J}_n$, we have

$$\lim_{n\to\infty} \frac{|J_n|}{\text{meas}(\mathbb{J}_n)} = 1,$$

and thus for sufficiently large n,

$$|J_n| \geq \frac{1}{2}\text{meas}(\mathbb{J}_n).$$

But from calculus we find that

$$\text{meas}(\mathbb{J}_n) = |\mathscr{S}_1|\Big(\prod_{s\in\mathscr{S}_1\setminus s_0} |\Delta y_s|\Big) n^{(|\mathscr{S}_1|-1)/2}. \quad \square$$

Lemma 3. *If* $\alpha \in J_n$, *then*

$$C_n(\alpha)\,\mathbf{a}^\alpha \sim \frac{1}{(2\pi n)^{(|\mathscr{S}_1|-1)/2}\prod_{s\in\mathscr{S}_1}\sqrt{a_s}} \exp\Big(-\sum_{s\in\mathscr{S}_1}\frac{y_s^2}{2a_s}\Big) \tag{2.6}$$

uniformly for $y_s \in [\bar{y}_s,\ \bar{y}_s + \Delta y_s]$ $(s \in \mathscr{S}_1\setminus s_0)$, *i.e., as* $n \to \infty$,

$$\sup_{\substack{y_s\in[\bar{y}_s,\ \bar{y}_s+\Delta y_s]\\ (s\in\mathscr{S}_1\setminus s_0)}} \left| C_n(\alpha)\,\mathbf{a}^\alpha \Big/ \left\{ \frac{1}{(2\pi n)^{(|\mathscr{S}_1|-1)/2}\prod_{s\in\mathscr{S}_1}\sqrt{a_s}} \exp\Big(-\sum_{s\in\mathscr{S}_1}\frac{y_s^2}{2a_s}\Big)\right\} - 1 \right| \to 0.$$

Proof. The proof depends on Stirling's formula

$$m! = \sqrt{2\pi m}\, e^{-m} m^m (1 + R(m)),$$

where $R(m) \to 0$ as $m \to \infty$. Since $\alpha_s = a_s n + y_s\sqrt{n} \to \infty$ as $n \to \infty$ for $s \in \mathscr{S}_1$, we have, by using the definition (2.2) and Stirling's formula,

$$\begin{aligned} C_n(\alpha)\,\mathbf{a}^\alpha &= \frac{\sqrt{2\pi n}\, e^{-n} n^n \mathbf{a}^\alpha (1 + R(n))}{(2\pi)^{|\mathscr{S}_1|/2}\Big(\prod_{s\in\mathscr{S}_1}\sqrt{\alpha_s}\Big) e^{-n}\alpha^\alpha \prod_{s\in\mathscr{S}_1}(1 + R(\alpha_s))} \\ &= \frac{1 + R(n)}{(2\pi n)^{(|\mathscr{S}_1|-1)/2}\prod_{s\in\mathscr{S}_1}\sqrt{a_s + y_s/\sqrt{n}}} \\ &\quad \times \frac{1}{\prod_{s\in\mathscr{S}_1}(1 + y_s/(a_s\sqrt{n}))^{a_s n + y_s\sqrt{n}} \prod_{s\in\mathscr{S}_1}(1 + R(\alpha_s))}, \end{aligned}$$

where the last equality follows from the fact that

$$\alpha^\alpha = \mathbf{a}^\alpha \prod_{s\in\mathscr{S}_1} n^{\alpha_s} \prod_{s\in\mathscr{S}_1}(1 + y_s/(a_s\sqrt{n}))^{\alpha_s} = \mathbf{a}^\alpha n^n \prod_{s\in\mathscr{S}_1}(1 + y_s/(a_s\sqrt{n}))^{a_s n + y_s\sqrt{n}}.$$

Here we have used (1.8) and $\sum_{s\in\mathscr{S}_1} y_s = 0$. In order to prove (2.6), we need to verify the following formulas:

$$\sqrt{a_s + \frac{y_s}{\sqrt{n}}} \sim \sqrt{a_s} \tag{2.7}$$

and

$$F(n) = \prod_{s\in\mathscr{S}_1}\Big(1 + \frac{y_s}{a_s\sqrt{n}}\Big)^{a_s n + y_s\sqrt{n}} \sim \exp\Big(\sum_{s\in\mathscr{S}_1}\frac{y_s^2}{2a_s}\Big). \tag{2.8}$$

The first, (2.7), is obvious. Now we prove (2.8). Since

$$\ln F(n) = \sum_{s\in\mathscr{S}_1} (a_s n + y_s\sqrt{n}\,) \ln\Big(1 + \frac{y_s}{a_s\sqrt{n}}\Big),$$

we find, by applying Taylor's formula to $\ln(1 + y_s/(a_s\sqrt{n}\,))$, that

$$\begin{aligned}\ln F(n) &= \sum_{s\in\mathscr{S}_1} (a_s n + y_s\sqrt{n}\,)\Big[\frac{y_s}{a_s\sqrt{n}} - \frac{1}{2}\Big(\frac{y_s}{a_s\sqrt{n}}\Big)^2 + \frac{1}{3}\Big(\frac{\tilde{y}_s}{a_s\sqrt{n}}\Big)^3\Big] \\ &= \sum_{s\in\mathscr{S}_1}\Big[y_s\sqrt{n} + \frac{{y_s}^2}{2a_s} + \frac{1}{a_s\sqrt{n}}\Big(-\frac{{y_s}^3}{2a_s} + \frac{{\tilde{y}_s}^3}{3{a_s}^2} + \frac{y_s{\tilde{y}_s}^3}{3\sqrt{n}{a_s}^3}\Big)\Big],\end{aligned}$$

where $\tilde{y}_s \in (0, y_s)$ and $y_s \in [\bar{y}_s, \bar{y}_s + \Delta y_s]$. Now using $\sum_{s\in\mathscr{S}_1} y_s = 0$, we obtain that for sufficiently large n

$$\ln F(n) = \sum_{s\in\mathscr{S}_1}\Big(\frac{{y_s}^2}{2a_s} + O\Big(\frac{1}{\sqrt{n}}\Big)\Big).$$

This verifies (2.8) and hence (2.6) is proved. □

Lemma 4. *If parameters $\bar{y}_s$ and Δy_s, defined in $\mathbb{J}_n$, are given by*

$$\bar{y}_s = \begin{cases} \dfrac{\sqrt{a_s}}{s - s_0}, & s - s_0 > 0, \\ \dfrac{2\sqrt{a_s}}{s - s_0}, & s - s_0 < 0, \end{cases} \qquad \Delta y_s = \frac{\sqrt{a_s}}{|s - s_0|} \quad (s \in \mathscr{S}_1\backslash s_0) \tag{2.9}$$

and **a** *satisfies* (1.7)–(1.9), *then for sufficiently large n,*

$$\min_{\alpha\in J_n} C_n(\alpha)\,\mathbf{a}^\alpha \geq \frac{0.5}{(2\pi n)^{(|\mathscr{S}_1|-1)/2}\prod_{s\in\mathscr{S}_1}\sqrt{a_s}} \exp\{-2(|\mathscr{S}_1| - 1)|\mathscr{S}_1|\}, \tag{2.10}$$

$$\min_{\alpha\in J_n}\Big|\sum_{s=-p}^{q} s\alpha_s + \lambda a n\Big| \geq \sqrt{n}\sum_{s\neq s_0}\sqrt{a_s}. \tag{2.11}$$

Proof. By using Lemma 3, we have for sufficiently large n,

$$C_n(\alpha)\,\mathbf{a}^\alpha \geq \frac{0.5}{(2\pi n)^{(|\mathscr{S}_1|-1)/2}\prod_{s\in\mathscr{S}_1}\sqrt{a_s}} \exp\Big(-\sum_{s\in\mathscr{S}_1}\frac{{y_s}^2}{2a_s}\Big). \tag{2.12}$$

Since $y_s \in [\bar{y}_s, \bar{y}_s + \Delta y_s]$ for $s \neq s_0$, we have, on account of (2.8),

$$\sum_{s\in\mathscr{S}_1}\frac{{y_s}^2}{2a_s} \leq \frac{{y_{s_0}}^2}{2a_{s_0}} + \sum_{s\in\mathscr{S}_1\backslash s_0}\frac{2}{(s - s_0)^2} \leq \frac{{y_{s_0}}^2}{2a_{s_0}} + 2(|\mathscr{S}_1| - 1). \tag{2.13}$$

On the other hand, from $\sum_{s\in\mathscr{S}_1} y_s = 0$ and $a_{s_0} = \max_s a_s$, we see that

$${y_{s_0}}^2 = \Big(\sum_{s\in\mathscr{S}_1\backslash s_0} y_s\Big)^2 \leq \Big(\sum_{s\in\mathscr{S}_1\backslash s_0}\frac{2\sqrt{a_s}}{|s - s_0|}\Big)^2 \leq 4a_{s_0}(|\mathscr{S}_1| - 1)^2,$$

or

$$\frac{y_{s_0}{}^2}{2a_{s_0}} \le 2(|\mathscr{S}_1| - 1)^2.$$

Substituting this into (2.13) gives

$$\sum_{s\in\mathscr{S}_1} \frac{y_s{}^2}{2a_s} \le 2(|\mathscr{S}_1| - 1)|\mathscr{S}_1|,$$

and combining this with (2.12) yields (2.10).

We now turn to (2.11). By using (2.4b), (1.9), and (2.9) , we have that

$$\left|\sum_s s\alpha_s + \lambda an\right| = \sqrt{n}\left|\sum_{s\in\mathscr{S}_1} sy_s\right| = \sqrt{n}\left|\sum_{s\in\mathscr{S}_1\setminus s_0} (s - s_0)y_s\right|$$
$$\ge \sqrt{n}\sum_{s\ne s_0}\sqrt{a_s}, \quad \forall\alpha \in J_n \subset \mathbb{J}_n.$$

This concludes the proof of (2.11). □

3. Proof of the main theorem

Proof of Theorem. By using (1.6) repeatedly for $n := 0, \dots, n-1$, we can express v_j^n in terms of the initial data v_j^0 as

$$v_j^n = \sum_{\alpha\in I_n} C_n(\alpha)\,\mathbf{a}^\alpha v^0_{j+\sum_s s\alpha_s}, \tag{3.1}$$

where $\sum_s$ is the sum over s from $-p$ to q. It is also known that the solution of (1.5) is of the form

$$u(x, t) = u_0(x - at). \tag{3.2}$$

Thus, we have

$$v_j^n - u(x, t_n) = \sum_{\alpha\in I_n} C_n(\alpha)\,\mathbf{a}^\alpha\left(v^0_{j+\sum_s s\alpha_s} - u_0(x - at_n)\right), \tag{3.3}$$

where we have used the equality (2.1). The upper error bound of (1.11) is a special case of the error estimate for scalar conservation laws [6], but the coefficient given here is more accurate. We will not present the proof here.

In order to prove the lower error bound of (1.11) , we only need to verify that the first inequality holds for the Riemann initial data

$$u_0(x) = \begin{cases} M/2 & \text{for } x > 0, \\ 0 & \text{for } x = 0, \\ -M/2 & \text{for } x < 0. \end{cases} \tag{3.4}$$

From (1.4) and (3.4), we see that

$$v^0_{j+\sum_s s\alpha_s} = u_0\left(x_j + \Delta x\sum_s s\alpha_s\right),$$

and hence

$$v^0_{j+\sum_s s\alpha_s} - u_0(x_j - at_n)$$
$$= \begin{cases} -M, & -\sum_s s\alpha_s > j > \lambda an, \\ -M/2, & -\sum_s s\alpha_s = j > \lambda an \text{ or } -\sum_s s\alpha_s > j = \lambda an, \\ 0, & j > (\lambda an) \vee \left(-\sum_s s\alpha_s\right) \text{ or } j < (\lambda an) \wedge \left(-\sum_s s\alpha_s\right), \\ M/2, & -\sum_s s\alpha_s = j < \lambda an \text{ or } -\sum_s s\alpha_s < j = \lambda an, \\ M, & -\sum_s s\alpha_s < j < \lambda an, \end{cases}$$

where $c \vee d = \max\{c, d\}$ and $c \wedge d = \min\{c, d\}$. Substituting this into (3.3) yields for $j \neq \lambda an$

(3.5)
$$v^n_j - u(x_j, t_n) = \sum_{\alpha \in I_n} C_n(\alpha)\, \mathbf{a}^\alpha (v^0_{j+\sum_s s\alpha_s} - u_0(x_j - at_n))$$
$$= \begin{cases} -M \sum_{\{\alpha \in I_n\} \cap \{j < -\sum_s s\alpha_s\}} C_n(\alpha)\, \mathbf{a}^\alpha \\ \qquad -M/2 \sum_{\{\alpha \in I_n\} \cap \{j = -\sum_s s\alpha_s\}} C_n(\alpha)\, \mathbf{a}^\alpha & \text{for } j > \lambda an, \\ M \sum_{\{\alpha \in I_n\} \cap \{j > -\sum_s s\alpha_s\}} C_n(\alpha)\, \mathbf{a}^\alpha \\ \qquad +M/2 \sum_{\{\alpha \in I_n\} \cap \{j = -\sum_s s\alpha_s\}} C_n(\alpha)\, \mathbf{a}^\alpha & \text{for } j < \lambda an. \end{cases}$$

For simplicity, we assume $t = t_n$ for some $n\,(= t/\Delta t)$. Since $u(x, t_n)$ is a two-piecewise constant function and $|v^n_j| \leq M/2$, we have

(3.6a)
$$\begin{aligned} \|v_{\Delta x}(\cdot, t) - u(\cdot, t)\|_{L^1(\mathbb{R})} &= \|v_{\Delta x}(\cdot, t_n) - u(\cdot, t_n)\|_{L^1(\mathbb{R})} \\ &\geq \Delta x \sum_j |v^n_j - u(x_j, t_n)| - M\Delta x \\ &\geq \Delta x \sum_{j \neq \lambda an} |v^n_j - u(x_j, t_n)| - M\Delta x. \end{aligned}$$

We divide the sum in the last term of (3.6a) into two parts,

(3.6b)
$$\sum_{j \neq \lambda an} |v^n_j - u(x_j, t_n)| = \mathrm{I}_+ + \mathrm{I}_-,$$

where

$$\mathrm{I}_+ = \sum_{j > \lambda an} |v^n_j - u(x_j, t_n)|$$

and

$$\mathrm{I}_- = \sum_{j < \lambda an} |v^n_j - u(x_j, t_n)|.$$

Substituting (3.5) into I_+ gives

$$\begin{aligned}
\mathrm{I}_+ &= M \sum_{j>\lambda an} \sum_{\{\alpha\in I_n\}\cap\{j<-\sum_s s\alpha_s\}} C_n(\alpha)\,\mathbf{a}^\alpha \\
&\quad + M/2 \sum_{j>\lambda an} \sum_{\{\alpha\in I_n\}\cap\{j=-\sum_s s\alpha_s\}} C_n(\alpha)\,\mathbf{a}^\alpha \\
&\geq M/2 \sum_{j>\lambda an} \sum_{\{\alpha\in I_n\}\cap\{j\leq-\sum_s s\alpha_s\}} C_n(\alpha)\,\mathbf{a}^\alpha \\
&= M/2 \sum_{\{\alpha\in I_n\}\cap\{\lambda an<-\sum_s s\alpha_s\}} \Big(-\sum_s s\alpha_s - [\lambda an]\Big) C_n(\alpha)\,\mathbf{a}^\alpha \\
&\geq M/2 \sum_{\{\alpha\in I_n\}\cap\{\lambda an<-\sum_s s\alpha_s\}} \Big(-\sum_s s\alpha_s - \lambda an\Big) C_n(\alpha)\,\mathbf{a}^\alpha,
\end{aligned}$$

where $[\eta]$ means the largest integer less than or equal to η. Similarly, we have

$$\mathrm{I}_- \geq M/2 \sum_{\{\alpha\in I_n\}\cap\{\lambda an>-\sum_s s\alpha_s\}} \Big(\sum_s s\alpha_s + \lambda an\Big) C_n(\alpha)\,\mathbf{a}^\alpha.$$

Adding I_+ and I_- yields

$$\begin{aligned}
\mathrm{I}_+ + \mathrm{I}_- &\geq M/2 \sum_{\{\alpha\in I_n\}\cap\{\lambda an\neq-\sum_s s\alpha_s\}} \Big|\sum_s s\alpha_s + \lambda an\Big| C_n(\alpha)\,\mathbf{a}^\alpha \\
&= M/2 \sum_{\alpha\in I_n} \Big|\sum_s s\alpha_s + \lambda an\Big| C_n(\alpha)\,\mathbf{a}^\alpha.
\end{aligned}$$

By the definitions (2.3) and (2.4b) we know that $J_n \subset I_n$ and, furthermore, assume that the parameters in $\mathbb{J}_n$ are given by (2.9), so that (2.10) and (2.11) hold. Then, on account of (2.5), we obtain

(3.7)

$$\begin{aligned}
\mathrm{I}_+ + \mathrm{I}_- &\geq M/2 \sum_{\alpha\in J_n} \Big\{\Big|\sum_s s\alpha_s + \lambda an\Big| C_n(\alpha)\,\mathbf{a}^\alpha\Big\} \\
&\geq \frac{M}{2}\,|J_n| \min_{\alpha\in J_n}\Big|\sum_s s\alpha_s + \lambda an\Big| \min_{\alpha\in J_n} C_n(\alpha)\,\mathbf{a}^\alpha \\
&\geq \frac{M}{2}\,\frac{\sqrt{|\mathscr{S}_1|}}{2}\Big(\prod_{s\in\mathscr{S}_1\setminus s_0}\frac{\sqrt{a_s}}{|s-s_0|}\Big) n^{(|\mathscr{S}_1|-1)/2}\sqrt{n}\sum_{s\neq s_0}\sqrt{a_s} \\
&\quad \times \frac{0,5}{(2\pi n)^{(|\mathscr{S}_1|-1)/2}\prod_{s\in\mathscr{S}_1}\sqrt{a_s}} \exp\{-2(|\mathscr{S}_1|-1)|\mathscr{S}_1|\} \\
&= \frac{\sqrt{|\mathscr{S}_1|}}{8}\Big(\prod_{s\in\mathscr{S}_1\setminus s_0}\frac{1}{|s-s_0|}\Big)\frac{\exp\{-2(|\mathscr{S}_1|-1)|\mathscr{S}_1|\}}{(2\pi)^{(|\mathscr{S}_1|-1)/2}\sqrt{a_{s_0}}} M\Big(\sum_{s\neq s_0}\sqrt{a_s}\Big)\sqrt{n}.
\end{aligned}$$

It follows from $|s - s_0| \le (p+q)$, $1 \le |\mathcal{S}_1| \le (p+q+1)$, and $a_{s_0} \le 1$, that (3.8)

$$\mathrm{I}_+ + \mathrm{I}_- \ge \frac{1}{8(p+q)^{(p+q)}(2\pi)^{(p+q)/2}} \exp\{-2(p+q)(p+q+1)\} M \left(\sum_{s \ne s_0} \sqrt{a_s} \right) \sqrt{n}$$

$$= 2\, c(p, q)\, M \left(\sum_{s \ne s_0} \sqrt{a_s} \right) \sqrt{n}.$$

We can see that $c(p, q) > 0$ is a constant which depends only on p and q. Since the Riemann initial data (3.4) satisfies $|u_0(\cdot)|_{\mathrm{BV}(\mathbb{R})} = M$, combining (3.6) and (3.8) yields the desired lower error bound, provided

$$\Delta x \le \left[c(p, q) \sum_{s \ne s_0} \sqrt{a_s} \right]^2 t/\lambda.$$

This completes the proof of the main theorem. □

Acknowledgments

We would like to thank the referee for helpful comments on the presentation of this material.

Bibliography

1. P. Brenner, V. Thomée, and L. B. Wahlbin, *Besov spaces and applications to difference methods for initial value problems* (A. Dold and B. Eckmann, eds.), Lecture Notes in Math., vol. 434, Springer-Verlag, Berlin and New York, 1975.
2. M. Crandall and A. Majda, *Monotone difference approximations to scalar conservation laws*, Math. Comp. **34** (1980), 1–21.
3. E. Engquist and S. Osher, *One-sided difference approximation for nonlinear conservation laws*, Math. Comp. **36** (1981), 321–351.
4. A. Harten, J. M. Hyman, and P. D. Lax, *On finite difference approximations and entropy conditions for shocks*, Comm. Pure Appl. Math. **29** (1976), 297–322.
5. G. W. Hedstrom, *The rate of convergence of some difference schemes*, SIAM J. Numer. Anal. **5** (1968), 363–406.
6. N. N. Kuznetsov, *On stable methods for solving non-linear first order partial differential equations in the class of discontinuous functions*, Topics in Numerical Analysis III (Proc. Roy. Irish Acad. Conf.) (J. J. H. Miller, ed.), Academic Press, London, 1977, pp. 183–197.
7. M. M. Loève, *Probability theory* I, Graduate Texts in Math., vol. 45, Springer-Verlag, 1977, pp. 22–23.
8. R. Sanders, *On convergence of monotone finite difference schemes with variable spatial differencing*, Math. Comp. **40** (1983), 91–106.

Department of Mathematics and Statistics, Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6
E-mail address: tangtao@cs.sfu.ca

Department of Mathematics, Peking University, Beijing 100871, People's Republic of China
E-mail address: tengzh@bepc2.ihep.ac.cn